

L15 ANSWER 55 OF 101 CA COPYRIGHT 2002 ACS

AN 105:4735 CA

TI Monoclonal T cell responses to two epitopes on a single immunogen controlled by two distinct genes

AU Lai, Chang Hai; Babu, Uma Mahesh; Turchin, Howard A.; Maurer, Paul H.

CS Dep. Biochem., Thomas Jefferson Univ., Philadelphia, PA, 19107, USA

SO J. Immunol. (1986), 136(10), 3799-804

CODEN: JOIMA3; ISSN: 0022-1767

DT Journal

LA English

AB The fine specificities of immune T cells were studied in a system in which the response to the antigen can involve 2 immune response (Ir) genes and 2 epitopes on a single synthetic polypeptide immunogen. The (BALB/c .times. SJL)F1 (H-2d .times. H-2s) mice can respond to the random terpolymer poly(Glu55,Lys36,Phe9) (GLPhe) through the H-2d-linked Ir gene (Ir-d) or through the complementing Ir gene (Ir-dxs), which controls the immune response to poly(Glu,Phe), epitopes that are present in GLPhe. Nine groups of monoclonal T cells were obtained from (H-2d .times. H-2s)F1 mice immunized with GLPhe. These groups were delineated by the differences in major histocompatibility complex (MHC)-restriction on antigen-presenting cells and the cross-reactions with poly(Glu60,Phe40) (GPhe) or poly(Glu51,Lys34,Tyr15) (GLT). A unique T cell line was discovered that can react to the 3 polymers (GLPhe, GLT, and GPhe) even though GLT and GPhe immune T cells do not normally show reciprocal cross-reactions. The monoclonal T cells retain helper activities in the Mishell-Dutton culture. Although the activation of T

Applicants: Alexander Gad and Dora Lis e;
Serial No.: 09/816,989
Filed: March 23, 2001
Exhibit 29

bot